

Aberystwyth University

Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica

Philippe, Morgane; Tison, Jean-Louis; Fjøsne, Karen; Hubbard, Bryn; Kjær, Helle A.; Lenaerts, Jan T. M.; Drews, Reinhard; Sheldon, Simon G.; De Bondt, Kevin; Claeys, Philippe; Pattyn, Frank

Published in:
Cryosphere

DOI:
[10.5194/tc-10-2501-2016](https://doi.org/10.5194/tc-10-2501-2016)

Publication date:
2016

Citation for published version (APA):

Philippe, M., Tison, J.-L., Fjøsne, K., Hubbard, B., Kjær, H. A., Lenaerts, J. T. M., Drews, R., Sheldon, S. G., De Bondt, K., Claeys, P., & Pattyn, F. (2016). Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica. *Cryosphere*, 10(5), 2501-2516. <https://doi.org/10.5194/tc-10-2501-2016>

Document License CC BY

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk



Supplement of

Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica

Morgane Philippe et al.

Correspondence to: Morgane Philippe (mophilip@ulb.ac.be)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Supplementary materials











Fig. S1. Full vertical profile of water stable isotopes with, from left to right: a grey and black band indicating sections of sampling for major ions at 10 cm and 5 cm resolution, respectively; water stable isotopes, taken at 5 cm resolution for the entire core; major ions, taken at 5 cm resolution for discrete sections; normalized ECM conductivity (0.05 m running mean, expressed as multiple of standard deviation, σ); annual layer boundaries in the youngest (Green) and the oldest (Blue) estimates (each colour transition indicates a boundary).



Fig. S2. Full vertical profile, as in Fig. S1 but split in 17 sections for more visibility.


























Table S1. Sites information and SMB values *no significant trend during the 20th century **short record: only recent periods are compared ***when only a stacked SMB change is given, SMB from individual ice cores are inferred from the stacked record as if it was the same trend for all ice cores. Ref : reference period. Numbers in italic are inferred from the trend given in the referenced paper

Site name	Latitude	Longitude	Elevation (m a.s.l.)	Reference period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	Recent period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	Most recent period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	% change (50a - ref)	% change (20a - ref) except**	Method	Study	
Siple Dome	-81.6530	-148.9980	620	1890-1994	120	1922-1991	118			-1.67%		Ice core	Kaspari et al., 2004	
ITASE00-5	-77.6830	-123.9950	1828	1716-2000	140	1922-1991	141			0.71%		Ice core	Kaspari et al., 2004	
ITAE99-1	-80.6200	-122.6300	1350	1724-1998	139	1922-1991	146			5.04%		Ice core	Kaspari et al., 2004	
ITASE00-4	-78.0830	-120.0800	1697	1799-2000	189	1922-1991	193			2.12%		Ice core	Kaspari et al., 2004	
GRIDS C	-80.0100	-119.4300	1530	1903-1995	112	1970-1995	108.35			-3.26%		Ice core	Kaspari et al., 2004	
GRIDS B	-79.4600	-118.0500	1603	1922-1995	150	1970-1995	149.37			-0.42%		Ice core	Kaspari et al., 2004	
GRIDS A	-78.7300	-116.3300	1740	1831-1995	235	1922-1991	234			-0.43%		Ice core	Kaspari et al., 2004	
ITASE00-1	-79.3830	-111.2390	1791	1653-2001	220	1922-1991	222			0.91%		Ice core	Kaspari et al., 2004	
ITASE01-2	-77.8430	-102.9100	1353	1890-2001	427	1922-1991	436			2.11%		Ice core	Kaspari et al., 2004	
ITASE01-3	-78.1200	-95.6460	1633	1859-2001	325	1922-1991	331			1.85%		Ice core	Kaspari et al., 2004	
ITASE01-5	-77.0590	-89.1370	1246	1780-2001	388	1922-1991	342			-11.86%		Ice core	Kaspari et al., 2004	
ITASE01-6	-76.0970	-89.0170	1232	**		1978-1990	395	1978-1999	392.6	-0.61%		Ice core	Kaspari et al., 2004	
Gomez	-73.5900	-70.3600	1400	1855-2006	720	1970s-2006	925	1997-2006	1100	28.47%	53%	Ice core	Thomas et al., 2008	
Dyer Plateau	-70.6700	-64.8900	2002	1790-1989	549	1969-1989	593			8.00%		Ice core	Raymond et al., 1996	
James Ross Island	-64.2200	-57.6800	1640	1847-1980	443	1964-1990	578			30.47%		Ice core	Aristarain et al., 2004	
R1	-78.3075	-46.2728	718	1816-1998	204	*	204			0.00%		Ice core	Mulvaney et al., 2002	
Berkner B25	-79.5700	-45.7200	890	1816-1956	131	1965-1994	141			7.63%		Ice core	Ruth et al., 2004	
A	-72.6500	-16.6333	60	**		1975-1989	380	1980-1989	350		-8%	Ice core	Isaksson & Melvold, 2002	
E	-73.6000	-12.4333	700	**		1932-1991	324	1980-1991	277		-15%	Ice core	Isaksson & Melvold, 2002; Isaksson et al., 1996	
B39	-71.4100	-9.9000	655	**		1935-2007	818	1987-2007	818		0.00%	Ice core	Fernandoy et al., 2010	
FB0704	-72.0600	-9.5600	760	**		1962-2007	489	1987-2007	489		0.00%	Ice core	Fernandoy et al., 2010	
BAS-depot	-77.0333	-9.5000	2176	1816-1997	71	1965-1997	71			0.00%		Ice core	Hofstede et al., 2004	
B04	-70.6200	-8.3700	35	1892-1981	362	±95 1960-1980	325			-10.22%			Schlosser & Oerter, 2002	
CV	-76.0000	-8.0500	2400	1816-1997	62	1965-1997	68	±2 1992-1997	70	9.68%	13%	Ice core	Karlof et al., 2005	
B38	-71.1600	-6.7000	690	**		1960-2007	1257	1987-2007	1257		0.00%	Ice core	Fernandoy et al., 2010	
FB0702	-71.5700	-6.6700	539	**		1959-2007	547	1987-2007	500		-9%	Ice core	Fernandoy et al., 2010	
FB9816	-75.0000	-3.5037	2740	1800-1997	47	±17 1950-1997	51.5***			9.57%		Ice core	Oerter et al., 2000	
B31	-75.5800	-3.4300	2669	1816-1997	58.4	1966-1989	59.8			2.40%		Ice core	Oerter et al., 2000	
H	-70.5000	-2.4500	53	**		1953-1993	480	1980-1993	425		-11%	Ice core	Isaksson & Melvold, 2002	
NUS08-2	-87.8500	-1.8000	2583	1815-2007/8	67.4	±2.6 1963-2007/8	63.4	±4.2		-5.93%		Ice core	Anschutz et al., 2011	
S32	-70.3100	-0.8000	53	**		1995-2009	339	±36	318		-6%	Ice core	Schlosser et al., 2014	
G3	-69.8230	-0.6120	57	**		1993-2009	295	±29	288		-2%	Ice core	Schlosser et al., 2014	
FB9815	-74.9492	-0.5055	2840	1801-1997	59	±24 1950-1997	65***			10.17%		Ice core	Oerter et al., 2000	
G4	-70.9020	-0.4020	60	**		1983-2009	330	±21	323		-2%	Ice core	Schlosser et al., 2014	
M2	-70.3160	-0.1090	73	**		1981-2009	315	±22	302		-4%	Ice core	Schlosser et al., 2014	
G5	-70.5450	-0.0410	82	**		1983-2009	298	±21	290		-3%	Ice core	Schlosser et al., 2014	
K	-70.7500	0.0000	53	**		1954-1996	254	1980-1996	250		0%	Ice core	Isaksson & Melvold, 2002	
SPS	-90.0000	0.0000	2850	1816-1956	76.5	1965-1994	84.8	±3.3 1992-1997	84.5	±8.9	10.85%	10%	Ice core and poles	Mosley & Thompson, 1999
B32	-75.0023	0.0070	2882	1816-1997	63	1966-1997	80			26.98%		Ice core	Oerter et al., 2000	
EPICA DML	-75.0020	0.0680	2774	1915-2008	73	1964-2008	73.1	±1.7		0.14%		Firn core and radar	Fujita et al., 2011	
FB9808	-74.7507	0.9998	2860	1801-1997	68	±22 1950-1997	74.5***			9.56%		Ice core	Oerter et al., 2000	
FB9809	-74.4992	1.9608	2843	1801-1997	89	±29 1950-1997	97.5***			9.55%		Ice core	Oerter et al., 2000	
EPICA (Amundsenisen)	-75.0000	2.0000	2900	1865-1965	78	1966-1991	76			-2.56%		Ice core	Isaksson et al., 1996	
G8	-70.4100	2.0100	58	**		1991-2009	282	±26	273		-3%	Ice core	Schlosser et al., 2014	

Site name	Latitude	Longitude	Elevation (m a.s.l.)	Reference period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	Recent period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	Most recent period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	% change (50a - ref)	% change (20a - ref) except**	Method	Study
FB9814	-75.0837	2.5017	2970	1801-1997	64	±21	1950-1997	71***		10.94%		Ice core	Oerter et al., 2000
C	-72.2583	2.8911	2400	1955-1996	119		1965-1996	123		3.36%		Ice core	Isaksson et al., 1999
D	-72.5083	3.0000	2610	1955-1996	112		1965-1996	116		3.57%		Ice core	Isaksson et al., 1999
DML08	-75.7528	3.2828	2971	1919-96	60	±19	*	60		0.00%		Ice core	Oerter et al., 1999
E	-72.6750	3.6628	2751	1955-1996	55		1965-1996	59		7.27%		Ice core	Isaksson et al., 1999
DML02	-74.9683	3.9185	3027	1919-95	59	±14	*	59		0.00%		Ice core	Oerter et al., 1999
FB9810	-74.6672	4.0017	2980	1801-1997	86	±29	1950-1997	94.5***		9.88%		Ice core	Oerter et al., 2000
F	-72.8583	4.3514	2840	1955-1996	23		1965-1996	24		4.35%		Ice core	Isaksson et al., 1999
S100	-70.2333	4.8000	48	1816-2000	292		1956-2000	284	1991-2000	-2.74%	-11%	Ice core	Kaczmarska et al., 2004
S20	-70.2417	4.8111	63	1955-1996	271		1965-1996	265		-2.21%		Ice core	Isaksson et al., 1999
FB0601	-75.2470	4.8440	3090	1915-2008	52		1964-2008	51.6	±1.2	-0.77%		Firn core and radar	Fujita et al., 2011
FB9813	-75.1673	5.0033	3100	1816-1997	48		1950-1997	53***		10.42%		Ice core	Oerter et al., 2000
G	-73.0417	5.0442	2929	1955-1996	28		1965-1996	30		7.14%		Ice core	Isaksson et al., 1999
FB9804	-75.2503	6.0000	2630	1801-1997	50	±16	1950-1997	55***		10.00%		Ice core	Oerter et al., 2000
H	-73.3917	6.4606	3074	1955-1996	44		1965-1996	46		4.55%		Ice core	Isaksson et al., 1999
B33	-75.1670	6.4985	3160	1816-1997	45.9		1966-1989	55		19.83%		Ice core	Oerter et al., 2000, Sommer et al., 2000
FB9811	-75.0840	6.5000	3160	1801-1997	58	±16	1950-1997	64***		10.34%		Ice core	Oerter et al., 2000
DML09	-75.9333	7.2130	3156	1897-1996	45	±12	*	45		0.00%		Ice core	Oerter et al., 1999
DML10	-75.2167	7.2130	3364	1900-96	47	±11	*	47		0.00%		Ice core	Oerter et al., 1999
DML04	-74.3990	7.2175	3179	1905-1996	53	±15	*	53		0.00%		Ice core	Oerter et al., 1999
I	-73.8008	7.9406	3174	1955-1996	52		1965-1996	53		1.92%		Ice core	Isaksson et al., 1999
NUS07-1	74.7200	7.9800	3174	1815–2007/8	52	±2	1963–2007/08	55.9	±3.9	7.50%		Ice core	Anschutz et al., 2009
Site I	-73.7167	7.9833	3174	1815-2007	52	±1.3	1963-2007	56	±4.7	7.69%	0%	Ice core	Anschutz et al., 2009
DML06	-75.0007	8.0053	3246	1899-1996	50	±14	*	50		0.00%		Ice core	Oerter et al., 1999
NUS08-6	-81.7000	8.5700	2447	1815-2007/8	39.2	±1.5	1963-2007/8	49.2	±3.4	25.51%		Ice core	Anschutz et al., 2011
J	-74.0417	9.4917	3268	1955-1996	44		1965-1996	45	±4	2.27%		Ice core	Isaksson et al., 1999
FB0603	-75.1170	9.7240	3300	1915-2008	41		1964-2008	38	±0.9	-7.32%		Firn core and radar	Fujita et al., 2011
K	-74.3583	11.1036	3341	1955-1996	45		1965-1996	41		-8.89%		Ice core	Isaksson et al., 1999
L	-74.6417	12.7908	3406	1955-1996	45		1965-1996	41		-8.89%		Ice core	Isaksson et al., 1999
A28	-74.8617	14.7420	3466	1915-2008	44		1964-2008	44.5	±1	1.14%		Firn core and radar	Fujita et al., 2011
MC	-75.0112	14.8865	3470.4	1816-1884	40		1955-2000	39		-2.50%	15%	Ice core	Karlof et al., 2005
MD	-74.9706	14.9567	3470.8	1816-1884	42		1955-2000	40		-4.76%	26%	Ice core	Karlof et al., 2005
M	-75.0000	14.9964	3470	1816-1884	41	±0.7	1955-2000	41	±0.5	0.00%	22%	Ice core	Karlof et al., 2005
M150	-74.9900	15.0000	3470	1816-1997	43		1965-1997	48.5		12.79%		Ice core	Hofstede et al., 2004
M	-74.9917	15.0017	3453	1955-1965	51		1965-1996	45		-11.76%		Ice core	Isaksson et al., 1999
MB	-75.0294	15.0435	3470.5	1816-1884	39		1955-2000	42		7.69%	18%	Ice core	Karlof et al., 2005
MA	-74.9887	15.1134	3470.4	1816-1884	42		1955-2000	42		0.00%	14%	Ice core	Karlof et al., 2005
NUS08-5	-82.6300	17.8700	2544	1815-2007/8	35	±0.8	1963-2007/8	37.6	±2.3	7.43%		Ice core	Anschutz et al., 2011
NUS08-4	-82.8167	18.9000	2552	1815-2007/8	36.7	±0.9	1963-2007/8	36.1	±2.1	-1.63%		Ice core	Anschutz et al., 2011
NUS08-3	-84.1300	22.0000	2625	1815-2007/8	40.1	±1	1963-2007/8	45.3	±3.1	12.97%		Ice core	Anschutz et al., 2011
A35	-76.0660	22.4590	3586	1915-2008	35		1964-2008	39.2	±0.9	12.00%		Firn core and radar	Fujita et al., 2011
NUS07-2	-76.0700	22.4700	3582	1815-2007/8	33	±0.7	1963-2007/8	28	±2	-15.15%		Ice core	Anschutz et al., 2011
MP	-75.8880	25.8340	3661	1286-2008	33.1	±1.0	1964-2008	38.7	±0.9	16.92%	27%	Firn core and radar	Fujita et al., 2011
NUS07-3	-77.0000	26.0500	3589	1815-2007/8	22	±0.5	1963-2007/8	23.7	±1.7	7.73%		Ice core	Anschutz et al., 2009
IC12	-70.2458	26.3349	450	1816-2012	480	±10	1955-2012	630	±20	31.25%	42%	Ice core	This paper
DK190	-76.7940	31.9000	3741	1286-2008	28.7	± 0.9					19%	Firn core and radar	Fujita et al., 2011
NUS07-4	-78.2167	32.8500	3595	1815-2007/8	19	±0.5	1963-2007/8	17.5	±1.2	-7.89%		Ice core	Anschutz et al., 2009
NUS07-5	-78.6500	35.6300	3619	1815-2007/8	24	±0.5	1963-2007/8	20.1	±1.4	-16.25%		Ice core	Anschutz et al., 2011

Site name	Latitude	Longitude	Elevation (m a.s.l.)	Reference period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	Recent period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	Most recent period	SMB (10 ⁻³ m w.e.) (kg m ⁻² a ⁻¹)	% change (50a - ref)	% change (20a - ref) except**	Method	Study		
DF	-77.3170	39.7030	3810	1816-2001	26.3	1964-2008	28.8	±0.7	1995-2006	27.3	±0.4	9.51%	4%	Ice core	Igarashi et al., 2011
YM85	-71.5800	40.6300	2246	1816-2002	140	1965-2002	135					-3.57%		Ice core	Takahashi et al., 2009
H72	-69.2047	41.0906	1214	1831-1998	311	1973-1998	307					-1.29%		Ice core and poles	Nishio et al., 2002
NUS07-6	-80.7833	44.8500	3672	1815-2007/8	22	1902-2007/8	21					-4.55%		Ice core	Anschutz et al., 2009
G15	-71.2000	45.9800	2544	1816-1964	86	1964-1984	116					34.88%		Ice core	Moore et al., 1991
NUS07-8	-84.1833	53.5333	3452	1815-2007/8	32	±1.2 1963-2007/8	30	±2.1				-6.25%		Ice core	Anschutz et al., 2009
NUS07-7	-82.0700	54.5500	3725	1815-2007/8	29.4	±0.6 1963-2007/8	26.1	±1.9				-11.22%		Ice core	Anschutz et al., 2011
DT217	-75.7167	76.8333	2800	**		1998-2008	12	±1.72	2005-2008	12			0%	Stake arrays	Ding et al., 2011
DT364	-78.3333	77.0000	3380	**		1999-2008	62	±0.14	2005-2008	72			16%	Stake arrays	Ding et al., 2011
DT401	-79.0200	77.0000	3760	1816-1999	19	1963-1999	24		1999–2005	25	±16	26.32%	32%	Ice core	Ren et al., 2010; Ding et al.,
DT001	-70.8300	77.0700	2325	1810-1959	131	1959-1996	131					0.00%		Ice core	Zhang et al., 2006
Dome A	-80.3667	77.3500	4093	**		2005-2008	19	±0.25	2008-2009	21			11%	Stake arrays	Ding et al., 2011
DomeA	-80.3600	77.3600	4092	1815-1998	23	1963-1998	23					0.00%		Ice core	Jiang et al., 2012
LGB65	-71.8500	77.9200	1850	1815-1996	131	1960-1996	131					0.00%		Ice core	Xiao et al., 2004
DT008	-72.1667	77.9333	2390	**		1998-2008	118	±0.30	2005-2008	80			-32%	Stake arrays	Ding et al., 2011
VOSTOK	-78.4500	106.8300	3488	1816-2010	20.6	±0.3 1955-2010	21.5	±0.5	1958-2010	20.8		4.37%	1%	Snow pits and poles	Ekaykin et al., 2004
DSS	-66.7697	112.8069	1370	1816-2000	680	1970-2009	750					10.29%		Ice core	Roberts et al., 2015
LAW DOME	-66.7700	112.9800	1370	1816-1966	687	1966-2005	742					8.01%		Ice core	Morgan et al., 1991; van Ommen & Morgan, 2010
DomeC	-75.1200	123.3100	3233	1816-1998	25.3	1965-1998	28.3		1996-1998	39		11.86%	54%	Ice core and poles	Frezzotti et al., 2005
D6 A	-75.4400	129.8100	3027	1816-1998	36	±1.8 1966-1998	29	±1.4	1998-2002	39		-19.44%	8%	Ice core and poles	Frezzotti et al., 2005
D66	-68.9400	136.9400	2333	1966-1864	196	1965-2001	213	±13	2001-2003	197		8.67%	1%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2013
D2 A	-75.6200	140.6300	2479	1816-1998	20	±1.0 1966-1998	31	±1.6	1998-2002	30		55.00%	50%	Ice core and poles	Frezzotti et al., 2005
GV1	-70.8700	141.3800	2244	1816-2001	114	1965-2001	117	±7	2001-2003	96		2.63%	-16%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2013
GV2	-71.7100	145.2600	2143	1816-2001	112	1965-2001	112	±7	2001-2003	92		0.00%	-18%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2013
MdPtA	-75.5300	145.8600	2454	1816-1998	36	±1.8 1966-1998	45	±2.7	1998-2010	47		25.00%	31%	Ice core and poles	Frezzotti et al., 2005
GV3	-72.6300	150.1700	2137	1816-2001	81	1965-2001	84	±5	2001-2003	73		3.70%	-10%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2013
M2 A	-74.8000	151.2700	2278	1816-1998	17	±0.8 1966-1998	15	±7.5	1998-2002	8.5		-11.76%	-50%	Ice core and poles	Frezzotti et al., 2005
GV4	-72.3900	154.4800	2126	1816–2001	119	1965–2001	100	±6	2001–2003	96		-15.97%	-19%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2013
31DPT A	-74.0300	155.9600	2069	1816-1998	98	±4.9 1966-1998	112	±5.6	1998-2002	98		14.29%	0%	Ice core and poles	Frezzotti et al., 2005
GPS2A	-74.6400	157.5020	1804	1816-1998	60	±3.0 1966-1998	54	±2.7	1993-2000	55		-10.00%	-8%	Ice core and poles	Frezzotti et al., 2005
GV5	-71.8900	158.5400	2184	1816-2001	129	1965-2001	129	±7	2001-2004	135		0.00%	5%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2007
GV7	-70.6800	158.8600	1947	1854-2001	237	1965-2001	241	±13	2001-2004	252		1.69%	6%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2007
Talos Dome	-72.7700	159.0800	2316	1816-2001	83.6	1966-1996	86.6		2001-2010	68		3.59%	-19%	Ice core and poles	Magand et al., 2004;Frezzotti et al., 2007; 2013
Hercules Neve	-73.1000	165.4000	2960	1816-1966	118	1966-1992	129					9.32%		Ice core	Stenni et al., 1999

References for Table S1

- Anschütz, H., Müller K., Isaksson, E., McConnell, J. R., Fischer, H., Miller, H., Albert, M., and Winther, J.-G.: Revisiting sites of the South Pole Queen Maud Land Traverses in East Antarctica: accumulation data from shallow firn cores, *J. Geophys. Res.*, 114, D24106, doi: 10.1029/2009JD012204, 2009.
- Ekaykin, A. A., Lipenkov, V. Y., Kuzmina, I., Petit, J. R., Masson-Delmotte, V., and Johnsen, S. J.: The changes in isotope composition and accumulation of snow at Vostok station, East Antarctica, over the past 200 years, *Ann. Glaciol.*, 39, 569–575, doi: 10.3189/172756404781814348, 2004.
- Frezzotti, M., Urbini, S., Proposito, M., Scarchilli, C., and Gandolfi, S.: Spatial and temporal variability of surface mass balance near Talos Dome, East Antarctica, *J. Geophys. Res.*, 112, F02032, doi: 10.1029/2006JF000638, 2007.
- Igarashi, M., Nakai, Y., Motizuki, Y., Takahashi, K., Motoyama, H., and Makishima, K.: Dating of the Dome Fuji shallow ice core based on a record of volcanic eruptions from AD 1260 to AD 2001, *Polar Sci.*, 5, 411–420, doi: 10.1016/j.polar.2011.08.001, 2011.
- Magand, O., Frezzotti, M., Pourchet, M., Stenni, B., Genoni, L., and Fily, M.: Climate variability along latitudinal and longitudinal transects in east Antarctica, *Ann. Glaciol.*, 39, 351–358, doi: 10.3189/172756404781813961, 2004.
- Morgan, V. I., Goodwin, I. D., Etheridge, D. M., and Wookey, C. W.: Evidence from Antarctic ice cores for recent increases in snow accumulation, *Nature*, 354(6348), 58–60, doi: 10.1038/354058a0, 1991.
- Mulvaney, R., Oerter, H., Peel, D. A., Graf, W., Arrowsmith, C., Pasteur, E. C., Knight, B., Littot, G. C., and Miners, W. D.: 1000-year ice core records from Berkner Island, Antarctic, *Ann. Glaciol.*, 35, 45–51, doi: 10.3189/172756402781817176, 2002.
- Raymond, C., Weertman, B., Thompson, L., Mosley-Thompson, E., Peel, D., and Mulvaney, R.: Geometry, motion and mass balance of Dyer Plateau, Antarctica, *J. Glaciol.*, 42, 510–518, doi: 10.3198/1996JoG42-142-510-518, 1996.
- Roberts, J., Plummer, C., Vance, T., van Ommen, T., Moy, A., Poynter, S., Treverrow, A., Curran, M., and George, S.: A 2000-year annual record of snow accumulation rates for Law Dome, East Antarctica, *Clim. Past*, 11, 697–707, doi: 10.5194/cp-11-697-2015, 2015.
- Ruth, U., Wagenbach, D., Mulvaney, R., Oerter, H., Graf, W., Pulz, H., and Littot, G.: Comprehensive 1000 year climate history from an intermediate depth ice core from the south dome of Berkner Island, Antarctica: methods, dating and first results, *Ann. Glaciol.*, 39, 146–154, doi: 10.3189/172756404781814104, 2004.
- Schlosser, E., Anschütz, H., Divine, D., Martma, T., Sinisalo, A., Altnau, S., and Isaksson, E.: Recent climate tendencies on an East Antarctic ice shelf inferred from a shallow firn core network, *J. Geophys. Res.*, 119, 6549–6562, doi: 10.1002/2013jd020818, 2014.
- Sommer, S., Appenzeller, C., Röthlisberger, R., Hutterli, M. A., Stauffer, B., Wagenbach, D., Oerter, H., Wilhelms, F., Miller, H., and Mulvaney, R.: Glacio-chemical study spanning the past 2 kyr on three ice cores from Dronning Maud Land, Antarctica, 1. Annually resolved accumulation rates, *J. Geophys. Res.*, 105, 29411–29421, doi: 10.1029/2000jd900449, 2000.
- Stenni, B., Caprioli, R., Cimmino, L., Cremisini, C., Flora, O., Gagnani, R., Longinelli, A., Maggi, V., and Torcini, S.: 200 years of isotope and chemical records in a firn core from Hercules Neve, northern Victoria Land, Antarctica, *Ann. Glaciol.*, 29, 106–112, doi: 10.3189/172756499781821175, 1999.
- Takahashi, H., Yokoyama, T., Igarashi, M., Motoyama, H., and Suzuki, K.: Resolution of environmental variation by detail analysis of YM85 shallow ice core in Antarctica, *Bull. Glaciol. Res.*, 27, 16–23, 2009.
- van Ommen, T. D., and Morgan, V.: Snowfall increase in coastal East Antarctica linked with southwest Western Australian drought, *Nat. Geosci.*, 3, 267–272, doi: 10.1038/ngeo761, 2010.
- Xiao, C., Mayewski, P. A., Qin, D., Li, Z., Zhang, M., and Yan, Y.: Sea level pressure variability over the southern Indian Ocean inferred from a glaciochemical record in Princess Elizabeth Land, east Antarctica, *J. Geophys. Res.*, 109, D16101, doi: 10.1029/2003JD004065, 2004.